arXiv:1208.6238v1 [math.FA] 30 Aug 2012

A NOTE ON THE POLYNOMIAL BOHNENBLUST–HILLE INEQUALITY

D. NUÑEZ-ALARCÓN*

Abstract. Recently, in paper published in the *Annals of Mathematics*, it was shown that the Bohnenblust–Hille inequality for (complex) homogeneous polynomials is hypercontractive. However, and to the best of our knowledge, there is no result providing (nontrivial) lower bounds for the optimal constants for n-homogeneous polynomials ($n > 2$). In this short note we provide lower bounds for these famous constants.

1. Introduction

The Bohnenblust–Hille inequality for complex homogeneous polynomials ([1], 1931) asserts that there is a function $D : \mathbb{N} \to [1, \infty)$ such that for every m-homogeneous polynomial P on $\mathbb{C}^N$, the $\ell_{\frac{2m}{m+1}}$-norm of the set of coefficients of P is bounded above by $D(m)$ times the supremum norm of P on the unit polydisc. In([2], 2011) it was proved that

$$D(m) \leq \left(1 + \frac{1}{m}\right)^{m-1} \sqrt{m} \left(\sqrt{2}\right)^{m-1} \tag{1}$$

which yields the hypercontractivity of the inequality.

The last few years experienced the rising of several works dedicated to estimating the Bohnenblust–Hille constants ([2, 3, 4, 6, 7]) and also unexpected connections with Quantum Information Theory appeared (see, e.g., [5]). There are in fact four cases to be investigated: polynomial (real and complex cases) and multilinear (real and complex cases). We can summarize in a sentence the main information from the recent preprints: the Bohnenblust–Hille constants are, in general, extraordinarily smaller than the first estimates predicted.

For example, now it is known that the Bohnenblust–Hille constants for the multilinear case behave in a subpolynomial way. In view of this, the investigation of lower bounds for the Bohnenblust–Hille constants seems to be an important task. The existing results for multilinear mappings and real scalars are highly nontrivial. For instance, in [4] it is shown that for multilinear mappings and real scalars one has that

$$C(m) \geq 2^{1-\frac{1}{m}}$$

and it is still open whether these estimates are sharp or not. Thus, the possibility of the boundedness of the multilinear Bohnenblust–Hille constants is open. In this short note we shall show that, if $m \geq 2$, then

$$D(m) \geq \frac{\left(2 + 2^m\right)^{\frac{m+1}{2m}}}{\sqrt{4 + 2^{m+1}}}.$$

2. The result

To the best of our knowledge, the only nontrivial lower bound for the constants of the (complex) polynomial Bohnenblust–Hille inequality is

$$D(2) \geq 1.1066,$$

proved in [6]. The lack of known estimates for the norms of complex polynomials of higher degrees was a barrier to obtain nontrivial lower estimates for $D(m)$ with $m > 2$. Our result provides nontrivial constants:

Key words and phrases. complex polynomials, multilinear operators, Bohnenblust–Hille inequality.

2010 Mathematics Subject Classification. 46G25, 12Y99.

*Supported by Capes.

Theorem 2.1. *For every* $m \geq 2$,

$$D(m) \geq \frac{(2+2^m)^{\frac{m+1}{2m}}}{\sqrt{4+2^{m+1}}} > 1.$$

Proof. Let $P_2 : \ell_\infty^2 \to \mathbb{C}$ be a 2-homogeneous polynomial given by

$$P_2(z_1, z_2) = az_1^2 + bz_2^2 + cz_1z_2.$$

with $a, b, c \in \mathbb{R}$ and $ab < 0$ and $|c(a+b)| \leq 4|ab|$. From [6] we know that

$$\|P_2\| = (|a| + |b|)\sqrt{1 + \frac{c^2}{4|ab|}}.$$

For each m, consider

$$P_m(z) = z_3 \ldots z_m P_2(z_1, z_2).$$

By choosing $a = -b = 1$, the $\ell_{\frac{2m}{m+1}}$ norm of the coefficients of P_m is

$$\left(\sqrt[m+1]{a^{2m}} + \sqrt[m+1]{b^{2m}} + \sqrt[m+1]{c^{2m}}\right)^{\frac{m+1}{2m}} = \left(2 + \sqrt[m+1]{c^{2m}}\right)^{\frac{m+1}{2m}}$$

and

$$\|P_m\| = \|P_2\| = (|a| + |b|)\sqrt{1 + \frac{c^2}{4\,|ab|}} = \sqrt{4 + c^2}.$$

Thus, we obtain

$$D(m) \geq f_m(x)$$

for all real numbers x, with

$$f_m(x) = \left(2 + \sqrt[m+1]{x^{2m}}\right)^{\frac{(m+1)}{2m}} . \left(\sqrt{4 + x^2}\right)^{-1}.$$

By solving $f_m'(x) = 0$ we conclude that $x = 2^{\frac{m+1}{2}}$ is a point where the maximum is achieved. Thus

$$D(m) \geq f_m(2^{\frac{m+1}{2}}) = (2 + 2^m)^{\frac{m+1}{2m}} . \left(\sqrt{4 + 2^{m+1}}\right)^{-1}.$$

A straightforward calculation shows that the last expression is always strictly greater than 1. □

References

[1] H.F. Bohnenblust and E. Hille, On the absolute convergence of Dirichlet series, Ann. of Math. (2) **32** (1931), 600–622.

[2] A. Defant, L. Frerick, J. Ortega-Cerdá, M. Ounaïes and K. Seip, The polynomial Bohnenblust–Hille inequality is hypercontractive, Ann. of Math. (2) **174** (2011), 485–497.

[3] D. Diniz, G.A. Muñoz-Fernández, D. Pellegrino and J.B. Seoane-Sepúlveda, The asymptotic growth of the constants in the Bohnenblust–Hille inequality is optimal, J. Funct. Anal. **263** (2012), 415–428.

[4] D. Diniz, G.A. Muñoz-Fernández, D. Pellegrino and J.B. Seoane-Sepúlveda, Lower bounds for the constants in the Bohnenblust–Hille inequality: the case of real scalars, Proc. Amer. Math. Soc., in press.

[5] A. Montanaro, Some applications of hypercontractive inequalities in quantum information theory, arXiv:1208.0161v2.

[6] G.A. Muñoz-Fernández, D. Pellegrino, J. Ramos Campos and J.B. Seoane-Sepúlveda, A geometric technique to generate lower estimates for the constants in the Bohnenblust–Hille inequalities, arXiv:1203.0793.

[7] D. Pellegrino and J.B. Seoane-Sepúlveda, New upper bounds for the constants in the Bohnenblust Hille inequality, J. Math. Anal. Appl. **386** (2012), 300–307.

Departamento de Matemática,
Universidade Federal da Paraíba,
58.051-900 - João Pessoa, Brazil.
E-mail address: `danielnunezal@gmail.com`